



HELAGO-CZ, s.r.o.
Commercial Register maintained by the Regional Court in Hradec
Králové
Section C, File 17879
Kladská 1082
500 03 Hradec Králové 3
Company ID: 25 96 39 61, VAT: CZ 25963961
Phone: 495 220 229, 495 220 394
Fax: 495 220 154
GSM gate: 602 123 096
E-mail: info@helago-cz.cz
Web: http://www.helago-cz.cz

MP1137 - Internal abdominal wall

Order code: **4003.MP1137**



Cena bez DPH

847,00 Eur

Price with VAT

1.024,87 Eur

Parameters

Quantitative unit

ks

This 3D model captures the internal surface of the anterior abdominal wall, a region oftentimes removed or damaged during dissection (and complementing our A8 abdominal specimen where the anterior wall has been removed). The parietal peritoneum has been removed from the internal surface of the specimen in order to more clearly demonstrate the relationships of the anterior abdominal muscle fibres and connective tissue structures as they converge on the midline. On the margins of the specimen, particularly superiorly, the horizontally-oriented transversus abdominus muscle fibres can be seen converging towards their aponeurosis (tendon sheet). In the inferior 1/3 of the model, we can see the termination of the posterior aspect of the aponeurosis forming the arcuate line. This marks the location where the aponeurosis changes its orientation relative to the rectus abdominus muscle (visible on either side of the midline); above the arcuate line the aponeurosis of the transversus abdominus muscle is evenly divided around the rectus abdominus muscle, while below the arcuate line all aponeurotic fibres pass anterior relative to the rectus abdominus. At this point, we can observe the inferior epigastric arteries (and accompanying veins) passing superiorly from their origins from the external iliac arteries and veins to pass into the anterior abdominal wall tissues. On the right side of the model we can appreciate how the orientation of the inferior epigastric artery relative to the fibres of the rectus abdominus muscle define the apex of the inguinal (Hesselbach's) triangle (missing only the base formed by the inguinal ligament, not present in this specimen). This region lateral to the inferior epigastric artery is a frequent site of direct hernias (which can be appreciated on the A8 abdomen model) given the relative weakness of the wall inferior to the arcuate line and lateral to the margin of the rectus abdominus muscle. In the midline, and dividing the two halves of the rectus abdominus muscle, is part of the median abdominal ligament - a draped fold of the parietal peritoneum that covers the urachus, a fibrous embryological remnant of the allantois, which extends from the bladder into the umbilical cord.