



Cena bez DPH	435,00 Eur
Price with VAT	526,35 Eur

Parameters

Quantitative unit	ks
-------------------	----

Clinical History

The patient was an 80-year old man who suddenly lost consciousness. On examination there was a right gaze palsy, a left hemiplegia and right hemiparesis.

Pathology

The specimens are coronal sections of the brain at the level of the mammillary bodies (specimen in which the cut surface of the brainstem where the cerebral peduncles and substantia nigra are also visible), and more anteriorly where part of both temporal lobes are included. A massive blood clot has replaced the cerebral tissue in the region of the left basal ganglia and internal capsule. The haemorrhage has originated in this area and has ruptured into the left lateral ventricle, and its temporal horn, destroying the walls of the left lateral ventricle and extending into adjacent brain tissue. The right lateral ventricle is also filled with blood, but its walls are intact. The blood clot forms a space-occupying lesion, which has expanded the left cerebral hemisphere, causing a shift of the mid-line structures to the right. There is subfalcine herniation of the left cingulate gyrus under the falx cerebri (the latter has been removed).

Note

Intra-cerebral haemorrhages of this type are usually a complication of systemic hypertension. The haemorrhage occurs due to rupture of a micro-aneurism of a branch of the striate arteries, which arise from the middle cerebral artery and supply the basal ganglia.