

HELAGO-CZ, s.r.o. Commercial Register maintained by the Regional Court in Hradec Králové Section C, File 17879 Kladská 1082 500 03 Hradec Králové 3 Company ID: 25 96 39 61, VAT: CZ 25963961 Phone: 495 220 229, 495 220 394 Fax: 495 220 154 GSM gate: 602 123 096 E-mail: info@helago-cz.cz Web: http://www.helago-cz.cz

MP2058 - Bronchopneumonia Order code: 4003.MP2058



Cena bez DPH Price with VAT

Parameters Quantitative unit 884,00 Eur 1.069,64 Eur

ks

## **Clinical History**

There is no clinical history for this specimen.

## Pathology

The specimen is a parasagittal section of the left lung. There are patchy regions of focal consolidations and discolorations caused by

congested and hyperaemic lung tissue distributed within both lobes; however, the upper lobe is more severely affected. The consolidation appears to be concentrated around the bronchioles, which are ectatic. The costal (pleural) surface of the upper lobe is especially discoloured.

## **Further Information**

Bronchopneumonia is a form of pneumonia characterized by inflammatory exudate within the intra-alveolar space resulting in consolidation that affects a large and continuous area of the lobe of a lung. It is one of the two anatomic classifications of pneumonia (the other being lobar pneumonia). The affected regions in this case shows classical focal red hepatization or consolidation in focal regions, which is due to vascular congestion with extravasation of red cells into alveolar spaces, along with increased numbers of neutrophils and fibrin. The filling of airspaces by the exudate leads to a gross appearance of solidification, or consolidation, of the alveolar parenchyma. Bronchopneumonia is a subtype of pneumonia. It is the acute inflammation of the bronchi, accompanied by inflamed patches in the peribronchial and peribronchiolar lobules of the lungs.

It is often contrasted with lobar pneumonia but in clinical practice the types are difficult to apply, as the patterns usually overlap. Bronchopneumonia (sometimes called lobular) often leads to lobar pneumonia as the infection progresses to affect an entire lobe. The same organism may cause one type of pneumonia in one patient, and another in a different patient. Causes Bronchopneumonia is usually a bacterial pneumonia rather than being caused by viral disease and is more commonly a hospital-acquired pneumonia than a communityacquired pneumonia, in contrast to lobar pneumonia.